

AD-NILPOTENT IDEALS AND THE SHI ARRANGEMENT

CHAO-PING DONG

ABSTRACT. We extend the Shi bijection from the Borel subalgebra case to the parabolic subalgebra case. In this process, there emerges naturally the I -deleted Shi arrangement $\text{Shi}(I)$, which interpolates between the Coxeter arrangement Cox and the Shi arrangement Shi , and which break the symmetry of Shi in certain symmetrical way. Among other things, we determine the characteristic polynomial $\chi(\text{Shi}(I), t)$ of $\text{Shi}(I)$ explicitly for A_{n-1} and C_n . More generally, let $\text{Shi}(G)$ be an arbitrary arrangement between Cox and Shi . Armstrong and Rhoades gave a formula on $\chi(\text{Shi}(G), t)$ for A_{n-1} recently. Inspired by this result, we obtain formulae on $\chi(\text{Shi}(G), t)$ for B_n , C_n and D_n .

1. INTRODUCTION

Let $\mathfrak{g}$ be a finite dimensional complex simple Lie algebra of rank l . Fix a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$. Then we have the root system $\Delta = \Delta(\mathfrak{g}, \mathfrak{h})$. Let V be the real vector space spanned by Δ . We denote by $\langle \cdot, \cdot \rangle$ the canonical inner product on V which is induced from $\mathfrak{h}$ of the Killing form of $\mathfrak{g}$. For convenience, we will equip V with an inner product $(\cdot, \cdot)$ which is a suitable scalar multiple of the canonical one. For any root $\alpha \in \Delta$, let $\mathfrak{g}_\alpha$ be the root space relative to α . Let $\Pi = \{\alpha_1, \dots, \alpha_l\} \subseteq \Delta^+$ be a fixed choice of simple and positive root systems of Δ , respectively. Let $\mathfrak{n} = \bigoplus_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$. Then $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}$ is a Borel subalgebra of $\mathfrak{g}$.

The abelian ideals of a Borel subalgebra were studied by Kostant [10, 11] in connection with the representation theory of semisimple Lie groups. In particular, D. Peterson's following theorem was detailed in [11]: there are 2^l abelian ideals of $\mathfrak{b}$, regardless of the type of $\mathfrak{g}$. Peterson's approach was to give a bijection between the abelian ideals of $\mathfrak{b}$ and a certain set of elements in the affine Weyl group $\widehat{W}$ of $\mathfrak{g}$. This surprising result led Cellini and Papi to find similarities for *ad-nilpotent* ideals of $\mathfrak{b}$, i.e., the ideals of $\mathfrak{b}$ which are contained in $\mathfrak{n}$. For example, they showed how to associate to any ad-nilpotent ideal $\mathfrak{i}$ of $\mathfrak{b}$ a uniquely determined element $w_i \in \widehat{W}$ in [5]. In [6], they gave further a bijection between the set of ad-nilpotent ideals of $\mathfrak{b}$ and the set of W -orbits in $\check{Q}/(h+1)\check{Q}$, where $\check{Q}$ is the coroot lattice and h is the Coxeter number.

On the other hand, we note that a bijection between the set of all ad-nilpotent ideals of $\mathfrak{b}$ and the dominant regions of the now-called Shi arrangement had been given by Shi in [17]. Thus, Theorems 3.2 and 3.6 there firstly count the number of all ad-nilpotent ideals of $\mathfrak{b}$. To state his result, let us recall some notations concerning hyperplane arrangement.

A *hyperplane arrangement* is a finite collection of affine hyperplanes in an Euclidean space. For example, the *Coxeter arrangement* associated with Δ^+ is the arrangement in V defined

2010 *Mathematics Subject Classification*. Primary 17B20, 05Exx.

Key words and phrases. ad-nilpotent ideal, Shi arrangement, quasi-antichain, characteristic polynomial.

by

$$(1) \quad \mathbf{Cox} := \{H_{\alpha,0} \mid \alpha \in \Delta^+\}.$$

Here for $\alpha \in \Delta^+$ and $k \in \mathbb{Z}$, we define a hyperplane

$$(2) \quad H_{\alpha,k} := \{v \in V \mid (v, \alpha) = k\}.$$

If $\mathcal{A}$ is a hyperplane arrangement of V , the connected components of $V - \bigcup_{H \in \mathcal{A}} H$ are called *regions*. For example, there are $|W|$ regions of $\mathbf{Cox}$, where W is the Weyl group associated with $\Delta(\mathfrak{g}, \mathfrak{h})$. For later use, let us single out the *dominant region* of $\mathbf{Cox}$ as follows:

$$(3) \quad V_\infty := \{v \in V \mid (v, \alpha) > 0, \forall \alpha \in \Delta^+\}.$$

Postnikov and Stanley introduced the idea of a *deformation* of the Coxeter arrangement in [12], which is an affine arrangement each of whose hyperplanes is parallel to some hyperplane of the Coxeter arrangement. The *Shi arrangement* $\mathbf{Shi}$ associated to Δ^+ can be viewed as such an example:

$$(4) \quad \mathbf{Shi} := \mathbf{Cox} \cup \{H_{\alpha,1} \mid \alpha \in \Delta^+\}.$$

This arrangement was defined by Shi in the study of the Kazhdan-Lusztig cellular structure of the affine Weyl group of type A , see Chapter 7 of [15]. A region of $\mathbf{Shi}$ is called *dominant* if it is contained in V_∞ .

For any $\mathfrak{h}$ -stable subset $\mathfrak{u}$ of $\mathfrak{g}$, let $\Phi_{\mathfrak{u}} \subset \Delta$ be the subset defined so that

$$\mathfrak{u} = \mathfrak{u} \cap \mathfrak{h} + \sum_{\alpha \in \Phi_{\mathfrak{u}}} \mathfrak{g}_\alpha.$$

Now let us cite the Shi bijection from Theorem 1.4 of [17] as follows:

Theorem 1. (Shi) *There exists a natural bijective map from the set of all the ad-nilpotent ideals of $\mathfrak{b}$ to the set of all the dominant regions of the hyperplane arrangement $\mathbf{Shi}$. The map sends $\mathfrak{i}$ to $\{v \in V_\infty \mid (v, \beta) > 1, \forall \beta \in \Phi_{\mathfrak{i}}; (v, \beta) < 1, \forall \beta \in \Delta^+ \setminus \Phi_{\mathfrak{i}}\}$.*

The first purpose of this paper is to generalize the Shi bijection from the Borel subalgebra case to the parabolic subalgebra case. Fix a subset $I \subseteq \Pi$. Let Δ_I be the sub root system of Δ spanned by I , and put $\Delta_I^+ = \Delta_I \cap \Delta^+$. Let

$$\mathfrak{p}_I = \mathfrak{h} + \sum_{\alpha \in \Delta_I \cup \Delta^+} \mathfrak{g}_\alpha$$

be the standard parabolic subalgebra of $\mathfrak{g}$ corresponding to I . Recall that an ideal $\mathfrak{i}$ of $\mathfrak{p}_I$ is called *ad-nilpotent* if for all $x \in \mathfrak{i}$, $\text{ad}_{\mathfrak{p}_I} x$ is nilpotent. Let

$$(5) \quad C_I = \{\beta \in \Delta^+ \setminus \Delta_I \mid \forall \alpha \in \Delta_I^+, \beta - \alpha \notin \Delta\}.$$

We define the *I-deleted Shi arrangement* as

$$(6) \quad \mathbf{Shi}(I) := \mathbf{Cox} \cup \{H_{\alpha,1} \mid \alpha \in C_I\}.$$

Again a region of $\mathbf{Shi}(I)$ is called dominant if it is contained in V_∞ . We endow Δ^+ with the usual partial ordering, i.e., $\alpha \leq \beta$ if $\beta - \alpha = \sum_{\gamma \in \Delta^+} c_\gamma \gamma$, where the c_γ are some non-negative integers. For any ad-nilpotent ideal $\mathfrak{i}$ of $\mathfrak{p}_I$, we put $\Psi_{\mathfrak{i}} = \cup_{\alpha} \{\beta \in C_I \mid \alpha \leq \beta\}$, where α runs over all the minimal elements of $(\Phi_{\mathfrak{i}}, \leq)$. Then we have

Theorem 2. *There exists a natural bijective map from the set of all the ad-nilpotent ideals of $\mathfrak{p}_I$ to the set of all the dominant regions of the hyperplane arrangement $\mathbf{Shi}(I)$. This map sends $\mathbf{i}$ to $\{v \in V_\infty \mid (v, \beta) > 1, \forall \beta \in \Psi_{\mathbf{i}}; (v, \beta) < 1, \forall \beta \in C_I \setminus \Psi_{\mathbf{i}}\}$.*

We note that the number of the ad-nilpotent ideals for $\mathfrak{p}_I$ was enumerated by Righi in Theorem 5.12 of [13] for $\mathfrak{g}$ classical, and in [14] for $\mathfrak{g}$ exceptional using GAP4. As a consequence, we also know the number of dominant regions of $\mathbf{Shi}(I)$.

Since $\mathbf{Shi}(\Pi)$ is $\mathbf{Cox}$ and $\mathbf{Shi}(\emptyset)$ is $\mathbf{Shi}$, the arrangement $\mathbf{Shi}(I)$ interpolates between the Coxeter arrangement and the Shi arrangement. More generally, one can consider

$$(7) \quad \mathbf{Shi}(G) = \mathbf{Cox} \cup \{H_{\alpha,1} \mid \alpha \in G\},$$

where G is any subset of Δ^+ .

The fundamental combinatorial object associated with a hyperplane arrangement $\mathcal{A}$ is its *intersection poset* $L(\mathcal{A})$, which is defined as the set of nonempty intersections of hyperplanes from $\mathcal{A}$, partially ordered by the *reverse* inclusion of subspaces. The *characteristic polynomial* $\chi(\mathcal{A}, t) \in \mathbb{Z}[t]$ of the arrangement $\mathcal{A}$ is defined by

$$(8) \quad \chi(\mathcal{A}, t) = \sum_{x \in L(\mathcal{A})} \mu(V, x) t^{\dim(x)},$$

where $\mu : L(\mathcal{A}) \times L(\mathcal{A}) \rightarrow \mathbb{Z}$ is the Möbius function of the poset $L(\mathcal{A})$, see (3.15) and section 3.11 of [18]. When $\mathfrak{g} = A_{n-1}$, Armstrong and Rhoades gave a formula for $\chi(\mathbf{Shi}(G), t)$ in [1] using the finite field method of Crapo and Rota [7]. This result is recalled in Theorem 4.3 below. Analyzing its proof, one sees that there are two key features: firstly, for a large prime p , the number $\chi(\mathbf{Shi}(G), p)$ is expressed as an alternating sum

$$(9) \quad \sum_{S \subseteq G} (-1)^{|S|} f(S),$$

where $f(S)$ is the number of vectors v in $\mathbb{F}_p^l - \mathbf{Cox}_p$ such that v satisfies $(H_{\alpha,1})_p$ for all $\alpha \in S$. Moreover, the summation is reduced to certain subsets of G , termed as the arc sets of G -partitions in [1]. Secondly, the number $f(S)$ is shown to be dependent only on $|S|$.

We find a uniform way to express the first feature. That is, as recorded in Lemma 3.7, for any finite dimensional complex simple Lie algebra $\mathfrak{g}$ of rank l , one can always express $\chi(\mathbf{Shi}(G), p)$ as in (9), and it suffices to take the summation over the *quasi-antichains* of Δ^+ contained in G . We refer the reader to Definition 3.6 for the term quasi-antichain. For the second feature, we find that the number $f(S)$ depends only on $|S|$ when $\mathfrak{g}$ is B_n or C_n . These lead us to Theorem 3 below. We note that although the second feature no longer holds for D_n , by a more careful analysis, we still obtain a formula for $\chi(\mathbf{Shi}(G), p)$ in this case, see Theorem 6.3.

Theorem 3. (Theorem 5.5 and Remark 5.6) *Let $\mathfrak{g}$ be B_n or C_n , $n \geq 2$. For any subset $G \subseteq \Delta^+$, let $\mathbf{Stir}(G, n-k)$ be the number of quasi-antichains of Δ^+ contained in G with length k . Then the characteristic polynomial of $\mathbf{Shi}(G)$ is given by*

$$\chi(\mathbf{Shi}(G), t) = \sum_{k=0}^n (-1)^k \mathbf{Stir}(G, n-k) \prod_{j=k}^{n-1} (t - 2j - 1).$$

Throughout this paper, we make the *convention* that a product equals to 1 whenever the lower index exceeds the upper index. For example, in the above theorem, the term $\prod_{j=k}^{n-1} (t - 2j - 1)$ is understood as 1 when $k = n$. Let us give some remarks about the number $\mathbf{Stir}(G, n - k)$. When $\mathfrak{g} = A_{n-1}$ and $G = \Delta^+$, it is nothing but $S(n, k)$, the number of partitions of $[n] := \{1, 2, \dots, n\}$ into k blocks, namely the *Stirling number of the second kind*, see the paragraph before (15) for more details. It has the recurrence

$$(10) \quad S(n, k) = kS(n - 1, k) + S(n - 1, k - 1),$$

see for example (1.93) of [18], and the combinatorial identity

$$(11) \quad \sum_{k=0}^n S(n, k)(t)_k = t^n,$$

where $(t)_k = t(t - 1) \cdots (t - k + 1)$, see for example (1.94d) of [18]. In Lemma 5.2, we find its analogs for C_n .

Going from $\mathbf{Shi}$ to $\mathbf{Shi}(G)$, the original symmetry of $\mathbf{Shi}$ is broken according to the choice of the subset $G \subseteq \Delta^+$. When G is arbitrarily chosen, we shall not expect $\mathbf{Shi}(G)$ to behave nicely. However, when we specialize G to be C_I , one gets the arrangement $\mathbf{Shi}(I)$, where the symmetry is broken kind of symmetrically. Thus we may expect $\mathbf{Shi}(I)$ to behave better. Indeed, when $\mathfrak{g} = A_{n-1}$, we find that: the polynomial $\chi(\mathbf{Shi}(I), t)$ factors into nonnegative integers and it depends only on $|I|$; moreover, the arrangement $\mathbf{Shi}(I)$ is inductively free in the sense of Athanasiadis [3], hence it is free in the sense of Terao [19], see Theorem 4.6. These results are based on the works of Athanasiadis [2, 3], Armstrong and Rhoades [1]. When $\mathfrak{g} = C_n$, we find that: $\chi(\mathbf{Shi}(I), t)$ always factors into nonnegative integers; moreover, if I contains $2e_n$, $\chi(\mathbf{Shi}(I), t)$ depends only on $|I|$, and the same conclusion holds if I does not contain $2e_n$, see Theorem 5.1. These results are obtained via Theorem 3 and Lemma 5.2.

The paper is organized as follows: we prove Theorem 2 in Section 2. We recall some preliminaries on the characteristic polynomial of a hyperplane arrangement in Section 3. In particular, the finite field method is described there. Moreover, we express $\chi(\mathbf{Shi}(G), p)$ as an alternating sum over the quasi-antichains in Lemma 3.7. Section 4 is devoted to the study of $\mathbf{Shi}(I)$ for A_{n-1} , while Section 5 handles the C_n case. Finally, a formula of $\chi(\mathbf{Shi}(G), t)$ is deduced for D_n in Section 6.

2. PROOF OF THEOREM 2

This section is devoted to the proof of Theorem 2. We begin with some preliminaries. Any subset P of Δ^+ inherits a partial ordering from $(\Delta^+, \leq)$. Let us denote the corresponding poset by $(P, \leq)$ or simply by P . Recall that a *dual order ideal* of P is a subset J of P such that if $t \in J$ and $t \leq s$ for $s \in P$, then $s \in J$. Recall also that an *antichain* of P is a subset of P consisting of pairwise non-comparable elements. Note that there is a canonical bijection between the dual order ideals of P and the antichains of P : given a dual order ideal, we send it to the set of its minimal elements; the inverse map sends the antichain $\{a_1, \dots, a_k\}$ to the dual order ideal which is the union of the principal dual order ideals $V_{a_1}, \dots, V_{a_k}$, where $V_a = \{b \in P \mid a \leq b\}$.

Theorem 2 will be proved by collecting the bijections in the following subsections.

2.1. Recall that a subset I of Π is fixed. Let us put

$$\mathcal{F}_I := \{\Phi \subseteq \Delta^+ \setminus \Delta_I : \text{if } \alpha \in \Phi, \beta \in \Delta^+ \cup \Delta_I \text{ and } \alpha + \beta \in \Delta^+, \text{ then } \alpha + \beta \in \Phi\}.$$

As noted in section 3 of [13], we have a bijection

$$(12) \quad \{\text{ad-nilpotent ideals of } \mathfrak{p}_I\} \rightarrow \mathcal{F}_I; \mathfrak{i} \mapsto \Phi_{\mathfrak{i}}.$$

2.2. For any $\Phi \in \mathcal{F}_I$, let $A(\Phi)$ be the set of all the minimal elements of $(\Phi, \leq)$. We note that $A(\Phi)$ is contained in C_I . Indeed, let β be any minimal element of $(\Phi, \leq)$, and take any $\alpha \in \Delta_I^+$, it suffices to show that $\beta - \alpha$ is not a root. Assuming the contrary gives $\beta + (-\alpha) \in \Delta^+ \setminus \Delta_I$. Since $\beta \in \Phi$, $-\alpha \in \Delta^+ \cup \Delta_I$ and $\Phi \in \mathcal{F}_I$, we conclude that $\beta - \alpha \in \Phi$, which contradicts to the minimality of β . Thus $A(\Phi)$ is an antichain of $(C_I, \leq)$ and we have a well-defined map

$$(13) \quad \mathcal{F}_I \rightarrow \{\text{antichains of } C_I\}; \Phi \mapsto A(\Phi).$$

Actually, the above map is bijective. To show this, it suffices to prove that for any antichain A of C_I , the set

$$\Phi(A) = \bigcup_{\beta \in A} \{\alpha \in \Delta^+ : \alpha \geq \beta\}$$

belongs to $\mathcal{F}_I$, which is precisely the content of Proposition 1.4 of [4] in view of the following

Lemma 2.1. *We have $C_I = \{\beta \in \Delta^+ \setminus \Delta_I \mid \forall \alpha \in I, \beta - \alpha \notin \Delta\}$.*

Proof. Fix any $\beta \in \Delta^+ \setminus \Delta_I$ such that $\beta - \alpha$ is not a root for any $\alpha \in I$, it suffices to show that $\beta - \gamma$ is not a root for any $\gamma \in \Delta_I^+$. Let us prove this by induction on the height of γ . There is nothing to prove when $\text{ht}(\gamma) = 1$. Suppose that we have proved it for all $\gamma' \in \Delta_I^+$ with $\text{ht}(\gamma') < r$. Now take $\gamma \in \Delta_I^+$ be such that $\text{ht}(\gamma) = r$. Choose $\alpha \in I$ such that $(\gamma, \alpha) > 0$. Then $\gamma - \alpha \in \Delta_I^+$. By assumption, $\beta - \alpha$ is not a root, thus $(\beta, \alpha) \leq 0$ and $(\beta - \gamma, \alpha) < 0$. Thus $\beta - \gamma$ is not a root since otherwise $\beta - (\gamma - \alpha)$ would be a root, contradicting to the induction hypothesis since $\text{ht}(\gamma - \alpha) = r - 1$. $\square$

2.3. We take the canonical bijection from $\{\text{antichains of } C_I\}$ to $\{\text{dual order ideals of } C_I\}$.

2.4. Given any dual order ideal Φ of C_I , define

$$R_\Phi = \{v \in V_\infty \mid (v, \beta) > 1, \forall \beta \in \Phi; (v, \beta) < 1, \forall \beta \in C_I \setminus \Phi\}.$$

Then the map

$$(14) \quad \{\text{dual order ideals of } C_I\} \rightarrow \{\text{dominant regions of } \text{Shi}(I)\}; \Phi \mapsto R_\Phi.$$

is well-defined in view of the following

Lemma 2.2. *The set R_Φ is non-empty. Hence it is a dominant region of $\text{Shi}(I)$.*

Proof. Let $\tilde{\Phi}$ be the unique dual order ideal of Δ^+ containing Φ , that is,

$$\tilde{\Phi} = \bigcup_{\beta \in \Phi} \{\alpha \in \Delta^+ : \alpha \geq \beta\}.$$

We claim that $C_I \setminus \Phi \subseteq \Delta^+ \setminus \tilde{\Phi}$. Indeed, if there exists $\alpha \in C_I \setminus \Phi$ such that $\alpha \in \tilde{\Phi}$, then $\alpha = \beta + \gamma$ for some $\beta \in \Phi$ and $\gamma \in \Delta^+$. Since $\alpha \in C_I$ and Φ is a dual order ideal of C_I , this

would imply that $\alpha \in \Phi$, which is absurd. Thus the claim follows, and we have $R_{\tilde{\Phi}} \subseteq R_{\Phi}$, where $R_{\tilde{\Phi}} = \{v \in V_{\infty} \mid (v, \beta) > 1, \forall \beta \in \tilde{\Phi}; (v, \beta) < 1, \forall \beta \in \Delta^+ \setminus \tilde{\Phi}\}$. Note that $R_{\tilde{\Phi}}$ is non-empty by Theorem 1. Thus, R_{Φ} is non-empty as well. Then it is immediate that R_{Φ} is a dominant region of $\text{Shi}(I)$. $\square$

Given any dominant region R of $\text{Shi}(I)$, we say that R is *above* the hyperplane $H_{\beta,1}$, $\beta \in \Delta^+$, if it contains R and the origin of V in different half spaces. Now define

$$\tau(R) = \{\beta \in C_I \mid R \text{ is above the hyperplane } H_{\beta,1}\}.$$

We note that $\tau(R)$ is a dual order ideal of C_I . Indeed, take any $\gamma \in C_I$ such that $\beta \leq \gamma$ for some $\beta \in \tau(R)$, then since R is contained in V_{∞} , we have

$$(v, \gamma) = (v, \beta) + (v, \gamma - \beta) \geq (v, \beta) > 1, \forall v \in R.$$

Therefore, $\gamma \in \tau(R)$ as desired. Now it is obvious that the map (14) is bijective with the inverse given by τ .

3. CHARACTERISTIC POLYNOMIAL AND THE FINITE FIELD METHOD

Let $\mathcal{A}$ be a collection of hyperplanes in the Euclidean space $\mathbb{R}^n$. Recall that its characteristic polynomial is defined in (8). In this section, we will collect some preliminaries concerning the determination of $\chi(\mathcal{A}, t)$. In particular, the finite field method due to Crapo and Rota [7] will be described. We will end up with a lemma which reduces $\chi(\text{Shi}(G), p)$ to an alternating sum over the quasi-antichains of Δ^+ contained in G .

3.1. Characteristic polynomial and Poincaré polynomial. The *Poincaré polynomial* of $\mathcal{A}$ is defined by

$$P(\mathcal{A}, t) = \sum_{x \in L(\mathcal{A})} \mu(V, x) (-t)^{n - \dim(x)}.$$

Comparing it with (8), one sees easily that

$$P(\mathcal{A}, t) = (-t)^n \chi(\mathcal{A}, -\frac{1}{t}) \text{ and } \chi(\mathcal{A}, t) = t^n P(\mathcal{A}, -\frac{1}{t}).$$

Example 3.1. Consider G_2 . Let $\Pi = \{\alpha_1, \alpha_2\}$, where $\alpha_1 = (0, 1)$ and $\alpha_2 = (\frac{\sqrt{3}}{2}, -\frac{3}{2})$. Then one can easily compute from definition that

- If $I = \{\alpha_1\}$, then $\chi(\text{Shi}(I), t) = (t - 3)(t - 5)$;
- If $I = \{\alpha_2\}$, then $\chi(\text{Shi}(I), t) = (t - 4)(t - 5)$. $\square$

Remark 3.2. In general, we shall not expect $\chi(\text{Shi}(I))$ to depend only on the cardinality of I . Indeed, since the negative of the coefficient of t in $\chi(\text{Shi}(I))$ is $|C_I| + |\Delta^+|$, one can easily construct examples such that $|I| = |I'|$ and $\chi(\text{Shi}(I), t) \neq \chi(\text{Shi}(I'), t)$.

Suppose that the normals to the hyperplanes of $\mathcal{A}$ span a subspace of $V \subseteq \mathbb{R}^n$ of dimension $r(\mathcal{A})$. We call $r(\mathcal{A})$ the *rank* of $\mathcal{A}$. We say a region of $\mathcal{A}$ is *relatively bounded* if its intersection with V is bounded. When $V = \mathbb{R}^n$, a region of $\mathcal{A}$ is relatively bounded if and only if it is bounded. A good reason to study the characteristic polynomial for hyperplane arrangements is given by the following classic theorem of Zaslavsky.

Theorem 3.3. (Section 2 of [20]) *For any hyperplane in $\mathbb{R}^n$, we have*

- The number of regions of $\mathcal{A}$ is $(-1)^n \chi(\mathcal{A}, -1)$;
- The number of relatively bounded regions of $\mathcal{A}$ is $(-1)^{r(\mathcal{A})} \chi(\mathcal{A}, 1)$.

Based on Shi's determination of the number of regions of **Shi** [15, 16], Headley obtained the characteristic polynomial of **Shi**.

Theorem 3.4. (Theorem 2.4 of [8]) *Let $\mathfrak{g}$ be a finite dimensional complex simple Lie algebra with rank l and Coxeter number h . Let V be the real vector space spanned by $\Delta(\mathfrak{g}, \mathfrak{h})$. Then for the Shi arrangement in V , we have*

$$P(\mathbf{Shi}, t) = (1 + ht)^l.$$

Recall that the Coxeter number $h = n + 1, 2n, 2n$ and $2(n - 1)$ for A_n, B_n, C_n and D_n respectively, see Section 3.18 of [9]. Denote the multiset of the roots of $\chi(\mathcal{A}, t)$ by $\exp(\mathcal{A})$, which are called the *exponents* of $\mathcal{A}$. For example, $\exp(\mathbf{Shi}) = \{h^l\}$ by the above theorem. Here and in what follows we write $\{a_1^{m_1}, a_2^{m_2}, \dots, a_r^{m_r}\}$ for a multiset, where $m_1, m_2, \dots, m_r$ denote multiplicities. The Factorization Theorem of Terao [19] states that the characteristic polynomial $\chi(\mathcal{A}, t)$ factors over the nonnegative integers for any free arrangement $\mathcal{A}$.

3.2. The finite field method. Athanasiadis [2] offered a different approach to Theorem 3.4 which did not rely on Shi's result. There the main tool was the *finite field method* of Crapo and Rota [7] which turned out to be very useful. Let us describe this method. Suppose that the defining equations for the hyperplanes in $\mathcal{A}$ have coefficients in $\mathbb{Z}$. Let $p \in \mathbb{Z}$ be a prime number and consider a hyperplane $H \in \mathcal{A}$ with defining equation $a_1x_1 + \dots + a_nx_n = b$, where $a_i, b \in \mathbb{Z}$. Then we define the following subset H_p of the finite vector space $\mathbb{F}_p^n$ by reducing the coefficients of H modulo p :

$$H_p := \{(x_1, \dots, x_n) \in \mathbb{F}_p^n \mid a_1x_1 + \dots + a_nx_n = b\}.$$

We note that when p is large enough, each H_p is a hyperplane in $\mathbb{F}_p^n$, and we call $\mathcal{A}_p := \{H_p \mid H \in \mathcal{A}\}$ the *reduced hyperplane arrangement* of $\mathcal{A}$.

Theorem 3.5. ([7]) *Let $p \in \mathbb{Z}$ be a large prime, and let $\mathcal{A}$ be a collection of hyperplanes in $\mathbb{R}^n$ whose hyperplanes have defining equations with coefficients in $\mathbb{Z}$. Then the characteristic polynomial of $\mathcal{A}$ satisfies*

$$\chi(\mathcal{A}, p) = \# \left(\mathbb{F}_p^n - \bigcup_{H \in \mathcal{A}} H_p \right).$$

That is, $\chi(\mathcal{A}, p)$ counts the number of points in the complement of the reduced arrangement $\mathcal{A}_p$ in the finite vector space $\mathbb{F}_p^n$.

A proof of the above theorem may also be found on pages 199–200 of [2].

3.3. Expressing $\chi(\mathbf{Shi}(G), p)$ as an alternating sum over the quasi-antichains. When $\mathfrak{g}$ is a finite dimensional simple Lie algebra over $\mathbb{C}$ with rank l , the arrangements **Shi** and **Cox**, thus any arrangement between them, can be realized as arrangements whose hyperplanes have defining equations in $\mathbb{Z}$. Let G be any subset of Δ^+ , recall from (7) that

$$\mathbf{Shi}(G) = \mathbf{Cox} \cup \{H_{\alpha, 1} \mid \alpha \in G\}.$$

We shall prepare a lemma for computing the characteristic polynomial of $\mathbf{Shi}(G)$. For this purpose, let us introduce

Definition 3.6. We call a subset $S \subseteq \Delta^+$ a *quasi-antichain* if for any two distinct elements α, β in S , the difference $\alpha - \beta$ is not a nonzero integer multiple of any root. We call the cardinality of a quasi-antichain its *length*.

Of course, any antichain of $(\Delta^+, \leq)$ is automatically a quasi-antichain of Δ^+ . The notion of quasi-antichain will be illustrated vividly in the following sections.

Lemma 3.7. *For the hyperplane arrangement $\mathbf{Shi}(G)$ defined in (7), we have*

$$\chi(\mathbf{Shi}(G), p) = \sum_{\substack{S \subseteq G \text{ is a quasi} \\ \text{-antichain of } \Delta^+}} (-1)^{|S|} f(S),$$

where p is a large prime, and for any subset $S \subseteq G$, $f(S)$ is the number of vectors v in $\mathbb{F}_p^l - \mathbf{Cox}_p$ such that v satisfies $(H_{\alpha,1})_p$ for all $\alpha \in S$.

Proof. By Theorem 3.5, it suffices to count the number of vectors in $\mathbb{F}_p^l - \mathbf{Shi}(G)_p$. By the principle of inclusion-exclusion (see for example Chapter 2 of [18]), we have

$$\chi(\mathbf{Shi}(G), p) = \sum_{S \subseteq G} (-1)^{|S|} f(S).$$

The lemma follows from the observation that $f(S) = 0$ if $S \subseteq G$ is not a quasi-antichain of Δ^+ . Indeed, in such a case, we can find two distinct roots α and β in S such that $\alpha - \beta = k\gamma$, where k is a positive integer and $\gamma \in \Delta^+$. Now if there exists v in $\mathbb{F}_p^l - \mathbf{Cox}_p$ satisfying the equations $(H_{\alpha,1})_p$ and $(H_{\beta,1})_p$, it satisfies the equation $(H_{k\gamma,0})_p = 0$ as well. Hence when p is large enough, it would be a solution of $(H_{\gamma,0})_p$, contradicting to the assumption that $v \notin \mathbf{Cox}_p$. $\square$

4. CHARACTERISTIC POLYNOMIAL OF $\mathbf{Shi}(I)$: TYPE A

Let $\mathfrak{g}$ be of type A_{n-1} , $n \geq 2$. We label the positive roots as $\alpha_{ij} = e_i - e_j$, $1 \leq i < j \leq n$. We will also refer to α_{ij} simply by ij . They span the real vector space $V = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 + \dots + x_n = 0\}$. The corresponding set of simple roots is $\Pi = \{12, 23, \dots, (n-1)n\}$. Recall from (2) that for us, the arrangements $\mathbf{Cox}$ and $\mathbf{Shi}$ etc are defined within V instead of $\mathbb{R}^n$. This section is mainly devoted to investigating the characteristic polynomial and the freeness of $\mathbf{Shi}(I)$.

4.1. Set partitions and quasi-antichains. Put $[n] := \{1, 2, \dots, n\}$. We say that $\pi = \{B_1, B_2, \dots, B_k\}$, where each B_i is a nonempty subset of $[n]$, is a *partition of $[n]$ into k blocks* if we have the disjoint union

$$[n] = B_1 \sqcup B_2 \sqcup \dots \sqcup B_k.$$

The *arc diagram* of π is drawn as follows: place the numbers $1, 2, \dots, n$ on a line and draw an arc between each pair $i < j$ such that

- i and j are in the same block of π ; and
- there is no $i < l < j$ such that i, l, j are in the same block of π .

A partition π of $[n]$ is called *nonnesting* if it does not contain arcs ij and kl such that $i < k < l < j$ —that is, no arc of π “nests” inside another. Figure 1 displays the arc diagrams for the partitions $\{\{1, 3\}, \{2\}, \{4, 5\}\}$ and $\{\{1, 5\}, \{2\}, \{3, 4\}\}$ of $[5]$. The first partition is nonnesting, while the second one is nesting since the arc 34 lies inside the arc 15 .

FIGURE 1. Two partitions of $[5]$

Let Γ_n be the staircase Young diagram. Following Shi [17], let us fill the positive roots of A_{n-1} into the boxes of Γ_{n-1} as follows: put the root ij into the $(n+1-j, i)$ -th box $B_{n+1-j,i}$. Figure 2 is an example with $n = 5$.

15	25	35	45
14	24	34	
13	23		
12			

FIGURE 2. The Young diagram Γ_4 for A_4

We will *identify* Δ^+ with Γ_{n-1} in this way, and transfer the partial ordering $\leq$ on Δ^+ to a partial ordering on Γ_{n-1} accordingly. We note that for any two positive roots α and β , $\alpha \leq \beta$ if and only if the box of β is to the north, or the west, or the northwest of the box of α . This describes the poset structure of Γ_{n-1} . By a bit abuse of notation, we will refer to this poset simply by Γ_{n-1} . We can characterize the quasi-antichains of Γ_{n-1} as follows: a subset S of Γ_{n-1} is a quasi-antichain if and only if in each row and column of Λ_n , there is at most one box of S . Now let us state an easy observation which should be well-known.

Lemma 4.1. *There exists a bijective map from the set of all the quasi-antichains of Γ_{n-1} to the set of all partitions of $[n]$. Moreover, this map sends the quasi-antichains of Γ_{n-1} with length $n - k$ to the partitions of $[n]$ with k blocks.*

Proof. Given any quasi-antichain A , then for each element $ij \in A$, draw an arc between the indices i and j . This procedure gives the arc graph of a partition $\pi(A)$. We then map A to

$\pi(A)$, which is easily seen to be bijective. Since a partition has k blocks if and only if its arc diagram has $n - k$ arcs, the second statement follows directly. $\square$

Remark 4.2. One sees easily that the map in the above lemma sends the set of all the antichains of Γ_{n-1} to the set of all the nonnesting partitions of $[n]$.

4.2. The characteristic polynomial and the freeness of $\text{Shi}(I)$. For any subset $G \subseteq \Gamma_{n-1}$, recall that

$$\text{Shi}(G) = \{x_i - x_j = 0, 1 \leq i < j \leq n\} \cup \{x_i - x_j = 1, ij \in G\}.$$

In [1], a partition π of $[n]$ is defined to be a G -partition if all its arcs are contained in G . Moreover, the G -Stirling number $\text{Stir}(G, k)$ is defined as the number of G -partitions with k blocks. The latter terminology is justified as follows: when $G = \Gamma_{n-1}$, $\text{Stir}(G, k)$ is nothing but $S(n, k)$, the number of partitions of $[n]$ with k blocks, which is a Stirling number of the second kind, see for example p. 73 of [18]. In view of Lemma 4.1, we have

$$(15) \quad \text{Stir}(G, k) = \#\{\text{quasi-antichains of } \Gamma_{n-1} \text{ contained in } G \text{ with length } n - k\}.$$

Now let us state a result of Armstrong and Rhoades in [1], which is deduced by the finite field method described in Theorem 3.5.

Theorem 4.3. (Theorem 3.2 of [1]) *For any subset $G \subseteq \Gamma_{n-1}$, the characteristic polynomial of $\text{Shi}(G)$ is given by*

$$\chi(\text{Shi}(G), t) = \sum_{k=0}^{n-1} (-1)^k \text{Stir}(G, n - k) \prod_{j=k+1}^{n-1} (t - j).$$

For our study, let us specialize G to be C_I , where I is some subset $I \subseteq \Pi$. The following result of Athanasiadis will be useful for determining the characteristic polynomial of $\text{Shi}(I)$ explicitly.

Theorem 4.4. (Theorem 2.2 of [3]) *Suppose that the subset $G \subseteq \Gamma_{n-1}$ has the following property: if $i < j < k$ and $ij \in G$, then $ik \in G$. Then*

$$\chi(\text{Shi}(G), t) = \prod_{1 < j \leq n} (t - c_j),$$

where $c_j = n + a_j - j + 1$ and $a_j = \#\{i < j \mid ij \in G\}$.

Remark 4.5. We note that the formulation of Corollary 3.3 in [1] is not entirely correct, as one may see from the A_2 case with $G = \{23\}$.

Theorem 4.6. *Let $\mathfrak{g}$ be A_{n-1} and let I be any subset of Π with cardinality $r \geq 1$. Then $\chi(\text{Shi}(I), t)$ is independent of the r elements that I contains, and*

$$(16) \quad \exp(\text{Shi}(I)) = \{(n - r)^{n-r}, (n - r + 1)^1, (n - r + 2)^1, \dots, (n - 1)^1\}.$$

In particular, the number of regions is $(n - r + 1)^{n-r-1} n! / (n - r)!$ and the number of bounded regions is $(n - r - 1)^{n-r-1} (n - 2)! / (n - r - 2)!$. Moreover, $\text{Shi}(I)$ is inductively free in the sense of Athanasiadis [3], hence it is free in the sense of Terao [19].

Proof. Fix any subset I of Π with cardinality $r \geq 1$. As noticed by Righi in §5.1 of [13], the poset $(C_I, \leq)$ is isomorphic to the poset Γ_{n-1-r} , see Figure 3 for two examples.




FIGURE 3. $\mathfrak{g} = A_4$, $I = \{45\}$, $I' = \{34\}$

Thus the number of quasi-antichains of Γ_{n-1} contained in C_I with length k equals to the number of quasi-antichains of Γ_{n-1-r} with length k , and equals to the number of partitions of $[n-r]$ into $n-r-k$ blocks by Lemma 4.1. As mentioned before (15), the latter number is $S(n-r, n-r-k)$, which is nonzero only if $0 \leq k \leq n-r$. Therefore by (15) and Theorem 4.3, we have

$$\chi(\text{Shi}(I), t) = \sum_{k=0}^{n-r} (-1)^k S(n-r, n-r-k) \prod_{j=k+1}^{n-1} (t-j).$$

This tells us that $\chi(\text{Shi}(I), t)$ depends only on r . To deduce its explicit expression, the first way is quoting the formula (11), which gives

$$\sum_{k=0}^{n-r} S(n-r, k)(t)_k = t^{n-r}.$$

Then after some elementary calculations, one arrives at (16). Alternatively, since $\chi(\text{Shi}(I), t)$ is independent of the r elements that I contains, we can focus on the special case that $I_0 = \{12, \dots, r(r+1)\}$. Then $C_{I_0} = \{ij \mid r+1 \leq i < j \leq n\}$, and applying Theorem 4.4 also gives (16).

For the last statement, we use Theorem 4.1 of [3]. That is, it suffices to rule out the following two possibilities:

- (a) there exists $1 \leq i < j < k \leq n$ such that $ij, jk \in C_I$ but $ik \notin C_I$;
- (b) there exists four distinct numbers $i_1 < j_1, i_2 < j_2 \in [n]$ such that $i_1 j_1, i_2 j_2$ gives *all* the edges between the vertices $\{i_1, j_1, i_2, j_2\}$ in C_I .

Suppose that (a) happens. Then by Lemma 2.1, the (not necessarily distinct) simple roots $i(i+1), (j-1)j, j(j+1), (k-1)k$ are not contained in I . Then $ik \notin C_I$ means that there is a simple root $\alpha \in I$ such that $\alpha_{ik} - \alpha$ is a root. Thus, α has to be $i(i+1)$ or $(k-1)k$, which is absurd. Now suppose that (b) happens. Without loss of generalities, we assume that $i_1 < i_2$. Then there are three cases: (1) $i_1 < j_1 < i_2 < j_2$; (2) $i_1 < i_2 < j_1 < j_2$; (3) $i_1 < i_2 < j_2 < j_1$.

Let us consider the last case, the other two cases are similar. Now by Lemma 2.1, the (not necessarily distinct) simple roots $i_1(i_1 + 1), (j_1 - 1)j_1, i_2(i_2 + 1), (j_2 - 1)j_2$ are not contained in I . Then one sees easily that $i_1j_2 \in C_I$, contradiction. $\square$

5. CHARACTERISTIC POLYNOMIAL OF $\text{Shi}(I)$: TYPE C

Let $\mathfrak{g}$ be of type C_n , $n \geq 2$. Then the positive roots can be chosen as $\Delta^+ = \{e_i - e_j \mid 1 \leq i < j \leq n\} \cup \{2e_i \mid 1 \leq i \leq n\}$. They span the real vector space $\mathbb{R}^n$. The corresponding set of simple roots is $\Pi = \{e_1 - e_2, e_2 - e_3, \dots, e_{n-1} - e_n, 2e_n\}$. This section is devoted to the proof of the following

Theorem 5.1. *Let $\mathfrak{g}$ be C_n and let I be any subset of Π with cardinality $1 \leq r \leq n - 1$. Then when I contains $2e_n$, $\chi(\text{Shi}(I), t)$ is independent of the other $r - 1$ simple roots that I contains, and we have*

$$(17) \quad \exp(\text{Shi}(I)) = \{(2n - 2r + 1)^{n-r+1}, (2n - 2r + 3)^1, (2n - 2r + 5)^1, \dots, (2n - 1)^1\}.$$

In particular, the number of regions is $(2n - 2r + 2)^{n-r}(2n)!!/(2n - 2r)!!$ and the number of bounded regions is $(2n - 2r)^{n-r}(2n - 2)!!/(2n - 2r - 2)!!$. Similarly, when I does not contain $2e_n$, $\chi(\text{Shi}(I), t)$ is independent of the r simple roots that I contains, and we have

$$(18) \quad \exp(\text{Shi}(I)) = \{(2n - 2r)^{n-r}, (2n - 2r + 1)^1, (2n - 2r + 3)^1, \dots, (2n - 1)^1\}.$$

In particular, the number of regions is $(2n - 2r + 1)^{n-r}(2n)!!/(2n - 2r)!!$ and the number of bounded regions is $(2n - 2r - 1)^{n-r}(2n - 2)!!/(2n - 2r - 2)!!$.

5.1. Quasi-antichains: characterization and representation. Following Shi [17], let us fill the positive roots of C_n into the staircase Young diagram Λ_n as follows: put $2e_i$ into the (i, i) -th box $B_{i,i}$, $1 \leq i \leq n$; put $e_i + e_j$ into the (i, j) -th box $B_{i,j}$, $1 \leq i < j \leq n$; put $e_i - e_j$ into the $(i, 2n + 1 - j)$ -th box $B_{i,2n+1-j}$, $1 \leq i < j \leq n$. The C_4 case is illustrated in Figure 4.

$2e_1$	$e_1 + e_2$	$e_1 + e_3$	$e_1 + e_4$	$e_1 - e_4$	$e_1 - e_3$	$e_1 - e_2$
	$2e_2$	$e_2 + e_3$	$e_2 + e_4$	$e_2 - e_4$	$e_2 - e_3$	
		$2e_3$	$e_3 + e_4$	$e_3 - e_4$		
			$2e_4$			

FIGURE 4. The staircase Young diagram Λ_4 for C_4

We will identify Δ^+ with Λ_n in this way, and transfer the partial ordering $\leq$ on Δ^+ to a partial ordering on Λ_n accordingly. We note that for any two positive roots α and β , $\alpha \leq \beta$ if and only if the box of β is to the north, or the west, or the northwest of the box of α . This describes the poset structure of Λ_n . By a bit abuse of notation, we will refer to this

poset simply by Λ_n . For later reference, we denote by T_{n-1} the subdiagram obtained from Λ_n by deleting the n -th column. The diagram T_{n-1} inherits a partial ordering from Λ_n . Let $L_1 = \{2e_i \mid 1 \leq i \leq n\}$, and define L_j , $2 \leq j \leq n-1$, to be the collection of roots on the j -th row and the j -th column of Λ_n . We can characterize the quasi-antichains of Λ_n as follows: a subset S of Λ_n is a quasi-antichain if and only if in each row and column of Λ_n , and in each L_j , $1 \leq j \leq n-1$, there is at most one box of S .

Put $\pm[n] = \{\pm 1, \dots, \pm n\}$. As in [2], we call the elements of $\pm[n]$ the *signed integers* from 1 to n . Let $S \subseteq \Delta^+$ be a quasi-antichain with cardinality $n - k$. Let us represent S in the following vivid way: put n boxes on a line and fill $-i, i$ in the i -th box for $1 \leq i \leq n$. If $2e_i \in S$ (there is at most one such i), delete the i -th box. If $e_i - e_j \in S$, for some $1 \leq i < j \leq n$, draw an arc between i and j from the below, and draw an arc between $-i$ and $-j$ from the above. If $e_i + e_j \in S$, for some $1 \leq i < j \leq n$, draw an arc between $-i$ and j from the above, and draw an arc between i and $-j$ from the below. Carrying out this process for all the $n - k$ roots in S , we call the resulting graph the *signed partition associated to S* and denote it by $\pi^\pm(S)$. For example, when $n = 5$ and $S = \{e_1 - e_2, e_2 - e_3\}$, the corresponding signed partition $\pi^\pm(S)$ is given in Figure 5.




FIGURE 5. A signed partition for C_5

We read $\pi^\pm(S)$ from the above in the following way: when there is an arc between two signed integers, single them out and join them with an arrow pointing to the right; when there is no arc on a box, single out the positive number in it. For example, reading Figure 5 from the above gives

$$-1 \rightarrow -2 \rightarrow -3, 4, 5.$$

Similarly, we read $\pi^\pm(S)$ from the below in the following way: when there is an arc between two signed integers, single them out, swap them, and then join them with an arrow pointing to the right; when there is no arc on a box, single out the negative number in it. For example, reading Figure 5 from the below gives

$$3 \rightarrow 2 \rightarrow 1, -4, -5.$$

We call $3 \rightarrow 2 \rightarrow 1$ the *negative* of $-1 \rightarrow -2 \rightarrow -3$, call -4 the negative of 4 etc. Then up to a choice of sign, there are 3 *essentially different* ordered parts for the signed partitions in Figure 5. One sees easily that similar things hold in general: no matter S contains a root of the form $2e_i$ or not, reading $\pi^\pm(S)$ from the above and the below always gives $2k$ ordered parts, where the negative of each part occurs exactly once, and up to a sign, there are k essentially different ordered parts of $\pi^\pm(S)$. The motivation of introducing $\pi^\pm(S)$ will be evident in the following subsection.

Let $S(\Lambda_n, n - k)$ be the number of quasi-antichains of Λ_n with length k , which is nonzero only if $0 \leq k \leq n$. Similarly, let $S(T_n, n - k)$ be the number of quasi-antichains of T_n with length k , which is nonzero only if $0 \leq k \leq n$. We prepare an elementary lemma concerning these two kind of numbers for later use.

Lemma 5.2. (i) *For Λ_n , we have the recurrence*

$$(19) \quad S(\Lambda_n, k) = S(\Lambda_{n-1}, k - 1) + (2k + 1)S(\Lambda_{n-1}, k)$$

and the identity

$$(20) \quad \sum_{k=0}^n S(\Lambda_n, k)[t]_{2k-1} = t^n,$$

where $[t]_{2k-1} = (t - 1)(t - 3) \cdots (t - 2k + 1)$.

(ii) *For T_n , we have the recurrence*

$$(21) \quad S(T_n, k) = S(T_{n-1}, k - 1) + (2k + 2)S(T_{n-1}, k)$$

and the identity

$$(22) \quad \sum_{k=0}^n S(T_n, k)[t]_{2k} = t^n,$$

where $[t]_{2k} = (t - 2)(t - 4) \cdots (t - 2k)$.

Proof. We only provide the proof for the Λ_n case. The T_n case is entirely similar. Since the subdiagram obtained from Λ_n by deleting the first row is isomorphic to Λ_{n-1} as posets, we denote it by Λ_{n-1} . A quasi-antichain S of Λ_n with length k can be formed in two ways: no element of S is chosen in the first row of Λ_n ; one element of S is chosen in the first row of Λ_n . The first number is $S(\Lambda_{n-1}, n - 1 - k)$. To figure out the second number, we note the following observation: for any box $B_{i,j}$ of Λ_n , where $i \geq 2$, thanks to the existence of $L_1, \dots, L_{n-1}$, there are *always* two elements in the first row of Λ_n which can not be chosen to form a quasi-antichain together with $B_{i,j}$. Suppose that we have an arbitrary quasi-antichain S' with length $k - 1$ of Λ_{n-1} at hand, and we want to add one element $B_{1,j}$ from the first row such that $S' \cup \{B_{1,j}\}$ is a quasi-antichain of Λ_n . Now for each element in S' , by the previous observation, there are two candidates in the first row that should be avoided. Moreover, there is no overlapping among these candidates due to the assumption that S' itself is a quasi-antichain. Thus there are exactly $2n + 1 - 2k$ allowable choices for $B_{1,j}$, and the second number is $(2n + 1 - 2k)S(\Lambda_{n-1}, n - k)$. Therefore, we have

$$(23) \quad S(\Lambda_n, n - k) = S(\Lambda_{n-1}, n - 1 - k) + (2n - 2k + 1)S(\Lambda_{n-1}, n - k),$$

which can be easily reformulated as (19). Finally, the identity (20) follows from (19) and an induction argument. $\square$

5.2. The formula for $f(S)$. Let S be a quasi-antichain of Δ^+ with length $n - k$. Let us adopt the way of Athanasiadis [2] to count $f(S)$. By definition, $f(S)$ counts the number of

n -tuples $(x_1, \dots, x_n) \in \mathbb{F}_p^n$ which are the solutions of $(H_{\alpha,1})_p$, for all $\alpha \in S$, and satisfy the following conditions:

$$\begin{aligned} x_i &\neq 0, & \text{for } 1 \leq i \leq n, \\ x_i \pm x_j &\neq 0, & \text{for } 1 \leq i < j \leq n. \end{aligned}$$

We think of such an n -tuple as a map from $\pm[n]$ to $\mathbb{F}_p$, sending i to the class $x_i \in \mathbb{F}_p$ and $-i$ to the class $-x_i$. We think of the elements of $\mathbb{F}_p$ as boxes arranged and labeled cyclically with the classes mod p . The top box is labeled with the zero class, the clockwise next box is labeled with the class 1 mod p etc. The n -tuples in $\mathbb{F}_p^n$ become placements of the signed integers from 1 to n in the p boxes, and $f(S)$ counts the number of placements which satisfy certain restrictions prescribed by the above conditions. To be more precise, our way of mapping $\pm[n]$ to $\mathbb{F}_p$ imposes the restriction that

(a) when a signed integer is placed in the class a , its negative is placed in the class $-a$;

The conditions $x_i \neq 0, x_i \pm x_j \neq 0$ impose the restrictions that

- (b) the zero class is always empty; and
- (c) distinct signed integers are placed in distinct classes.

Moreover, the n -tuple should satisfy $(H_{\alpha,1})_p$ for all $\alpha \in S$.

Lemma 5.3. *Let S be a quasi-antichain of Δ^+ with length $n - k$. Let p be a large prime, then*

$$f(S) = \frac{(p - 2n + 2k - 1)!!}{(p - 2n - 1)!!}.$$

Proof. For convenience, we call the classes from 1 to $(p-1)/2$, both included, the right half of the circle. Similarly, we call the classes from $(p+1)/2$ to $p-1$, both included, the left half of the circle. By definition of $\pi^\pm(S)$, besides some additional restrictions which will be studied case by case below, we should place its $2k$ ordered parts on the circle such that the restrictions (a-c) are met. Namely, we should always place the $2k$ ordered parts of $\pi^\pm(S)$ on the circle such that

- each ordered part should be placed consecutively and clockwise on the circle, and there should be no overlapping among them;
- each ordered part should be placed entirely on the right half, or entirely on the left half; if an ordered part is placed on the right half, then its negative should be placed on the left half according to (a).

Since an allowable placement of the right half gives an allowable placement of the left half, we can focus on what happens only on the right half. Thus, it boils down to put the k essentially different ordered parts there, with each part a choice of sign. Now let us discuss the further restrictions case by case.

Firstly, let us consider the case that $2e_i \in S$ for some $i \in [n]$. Then $-i$ is sent to the $(p-1)/2$ class, and i to the $(p+1)/2$ class. Since there are $(p-1)/2 - n$ empty boxes on the right half circle in total, the allowable ways to place the k essentially different parts there is $f(S) = 2^k \prod_{j=1}^k ((p-1)/2 - n + j)$, which equals to $(p - 2n + 2k - 1)!! / (p - 2n - 1)!!$, as desired.

Secondly, let us consider the case that there is no $i \in [n]$ such that $2e_i \in S$. There are two possibilities:

- (1) the class $(p-1)/2$ is empty. Therefore, to place the k essentially different parts in the right half circle, we should always avoid the class $(p-1)/2$. Since besides the class $(p-1)/2$, there are $(p-1)/2 - n - 1$ empty boxes there in total, the allowable ways are easily seen to be $2^k \prod_{j=1}^k ((p-1)/2 - n - 1 + j)$.
- (2) the class $(p-1)/2$ is filled. Then there must be an ordered part of $\pi^\pm(S)$ which is placed entirely on the right half circle such that the corresponding consecutive part ends with the class $(p-1)/2$. We have $2k$ different ways to choose this ordered part, which will then be placed in a unique way. Now we have to place the remaining $k-1$ essentially different ordered parts in the remaining classes of the right half. Since there is a choice of sign for each of these $k-1$ parts, and there are $(p-1)/2 - n$ empty boxes on the right half in total, the allowable ways are $(2k)2^{k-1} \prod_{j=1}^{k-1} ((p-1)/2 - n + j)$.

Summing the numbers in (1) and (2) also gives the desired expression for $f(S)$. $\square$

Let us record an interesting corollary of Lemma 5.3.

Corollary 5.4. *Let p be a large prime. For any subset $S \subseteq \Lambda_n$ with cardinality $0 \leq k \leq n$, the following are equivalent:*

- (i) S is a quasi-antichain of Λ_n ;
- (ii) in each row and column of Λ_n , and in each L_j , $1 \leq j \leq n-1$, there is at most one box of S ;
- (iii) $f(S)$ is nonzero;
- (iv) $f(S) = (p-2k-1)!!/(p-2n-1)!!$.

Let G be any subset of Δ^+ . In accordance with (15), we define

$$(24) \quad \text{Stir}(G, k) = \#\left\{ \text{quasi-antichains of } \Lambda_n \text{ contained in } G \text{ with length } n-k \right\}.$$

Theorem 5.5. *Let $\mathfrak{g}$ be C_n , $n \geq 2$. Then for any subset $G \subseteq \Lambda_n$, the characteristic polynomial of $\text{Shi}(G)$ is given by*

$$\chi(\text{Shi}(G), t) = \sum_{k=0}^n (-1)^k \text{Stir}(G, n-k) \prod_{j=k}^{n-1} (t-2j-1).$$

Proof. This follows from Lemma 3.7 and Lemma 5.3. $\square$

Remark 5.6. Since the positive roots for B_n are $\{e_i - e_j \mid 1 \leq i < j \leq n\} \cup \{e_i \mid 1 \leq i \leq n\}$, one sees immediately that Lemma 5.3, thus the theorem above, holds for B_n .

Example 5.7. When G is the empty set, by Theorem 5.5, we have

$$\chi(\text{Cox}, p) = (p-1)!!/(p-2n-1)!! = (p-1)(p-3) \cdots (p-2n+1).$$

When $G = \Gamma_{n-1}$, by Theorem 5.5, we have

$$\begin{aligned}
\chi(\text{Shi}, p) &= \sum_{k=0}^n (-1)^k S(\Lambda_n, n-k)(p-2k-1)(p-2k-3) \cdots (p-2n+1) \\
&= (-1)^n \sum_{k=0}^n S(\Lambda_n, n-k)(-p+2n-1) \cdots (-p+2k+3)(-p+2k+1) \\
&= (-1)^n \sum_{k=0}^n S(\Lambda_n, n-k)[-p+2n]_{2n-2k-1} \\
&= (-1)^n \sum_{k=0}^n S(\Lambda_n, k)[-p+2n]_{2k-1} \\
&= (-1)^n (-p+2n)^n \\
&= (p-2n)^n,
\end{aligned}$$

where in the next-to-last step we use the identity (20). $\square$

5.3. Proof of Theorem 5.1 when I contains $2e_n$. Fix any subset $I \subseteq \Pi$ with cardinality $1 \leq r \leq n-1$ which contains $2e_n$. As noticed by Righi in §5.2 of [13], the poset $(C_I, \leq)$ is isomorphic to the poset T_{n-r} , see Figure 6 for two examples.




FIGURE 6. $\mathfrak{g} = C_4$, $I = \{e_3 - e_4, 2e_4\}$, $I' = \{e_2 - e_3, 2e_4\}$

Thus the number of quasi-antichains of Λ_n contained in G with length k equals to the number of quasi-antichains of T_{n-r} with length k . Note that the latter number is $S(T_{n-r}, n-r-k)$, which is nonzero only if $0 \leq k \leq n-r$. Now by (24) and Theorem 5.5, we have

$$\chi(\text{Shi}(I), t) = \sum_{k=0}^{n-r} (-1)^k S(T_{n-r}, n-r-k) \prod_{j=k}^{n-1} (t-2j-1).$$

This tells us that $\chi(\text{Shi}(I), t)$ is independent of the other $r-1$ simple roots that I contains. To arrive at (17), after some elementary calculations similar to those in Example 5.7, it boils down to show

$$(25) \quad \sum_{k=0}^{n-r} S(T_{n-r}, k)[t]_{2k} = t^{n-r},$$

which is an easy consequence of Lemma 5.2(ii).

5.4. Proof of Theorem 5.1 when I does not contain $2e_n$. Fix any subset $I \subseteq \Pi$ with cardinality $1 \leq r \leq n-1$ which does not contain $2e_n$. This case is similar to the previous one. The only difference lies in that, as noticed by Righi in §5.2 of [13], the poset $(C_I, \leq)$ is now isomorphic to the poset Λ_{n-r} . Then replacing T_{n-r} in the previous case by Λ_{n-r} and quoting Lemma 5.2(i) instead finish the proof. $\square$

6. CHARACTERISTIC POLYNOMIAL OF $\text{ShI}(G)$: TYPE D

Let $\mathfrak{g} = D_n$, $n \geq 4$. We can choose $\Delta^+ = \{e_i \pm e_j \mid 1 \leq i < j \leq n\}$, then the corresponding set of simple roots is $\Pi = \{e_1 - e_2, e_2 - e_3, \dots, e_{n-1} - e_n, e_{n-1} + e_n\}$. They span the real vector space $\mathbb{R}^n$.

In this section, we will give a formula for $\chi(\text{ShI}(G), t)$, where G is an arbitrary subset of Δ^+ . As hinted by Lemma 3.7, this can be achieved by finding a formula for $f(S)$, where S is an arbitrary quasi-antichain of Δ^+ contained in G . In the A_{n-1} , B_n , C_n cases, we always have the nice property that $f(S)$ depends only on $|S|$. Unfortunately, as suggested by the following example, this is no longer true for D_n .

Example 6.1. Let us consider D_4 . Then

- $f(S) = (p-4)(p-5)$ when $S = \{e_1 + e_3, e_2 - e_3\}$;
- $f(S) = (p-3)(p-5)$ when $S = \{e_1 + e_2, e_3 + e_4\}$.

$\square$

However, by analyzing what happens in Example 6.1 more carefully, we can still obtain a formula for $f(S)$. Let $S \subseteq \Delta^+$ be a quasi-antichain with length $n-k$. Similar to the C_n case, we associate to S a signed partition $\pi^\pm(S)$. Reading $\pi^\pm(S)$ from the above and the below gives $2k$ ordered parts, where the negative of each part occurs exactly once. Thus up to a sign, there are k essentially different ordered parts. Let $n_1(S)$ be one half of the number of ordered parts with length 1 in $\pi^\pm(S)$. It is necessarily a nonnegative integer.

Similar to §5.2, we adopt the way of Athanasiadis [2] to count $f(S)$. By definition, $f(S)$ counts the number of n -tuples $(x_1, \dots, x_n) \in \mathbb{F}_p^n$ which are the solutions of $(H_{\alpha,1})_p$, for all $\alpha \in S$, and satisfy the conditions: $x_i \pm x_j \neq 0$, for $1 \leq i < j \leq n$. We think of such an n -tuple as a map from $\pm[n]$ to $\mathbb{F}_p$, sending i to the class $x_i \in \mathbb{F}_p$ and $-i$ to the class $-x_i$. We think of the elements of $\mathbb{F}_p$ as boxes arranged and labeled cyclically with the classes mod p . The top box is labeled with the zero class, the clockwise next box is labeled with the class 1 mod p etc. The n -tuples in $\mathbb{F}_p^n$ become placements of the signed integers from 1 to n in the p boxes, and $f(S)$ counts the number of placements which satisfy the following restrictions:

- (a) when a signed integer is placed in the class a , its negative is placed in the class $-a$;
- (b) there is at most one signed integers placed in each nonzero class;
- (c) there exists at most one $i \in [n]$ such that both i and $-i$ are placed in the zero class.

Moreover, the n -tuple should satisfy $(H_{\alpha,1})_p$ for all $\alpha \in S$.

Lemma 6.2. *Let S be a quasi-antichain of Δ^+ with length $n-k$. Let p be a large prime, then*

$$f(S) = \frac{(p-2n+2k+1)!!}{(p-2n+1)!!} - n_1(S) \frac{(p-2n+2k-1)!!}{(p-2n+1)!!},$$

where $n_1(S)$ is one half of the number of ordered parts with length 1 in $\pi^\pm(S)$.

Proof. By definition of $\pi^\pm(S)$, it suffices to count the placements of its $2k$ ordered parts on the circle such that the restrictions (a-c) are met. For convenience, we call the zero class combined with the right half the *extended* right half. Similarly, we call the zero class combined with the left half the *extended* left half. Therefore, we should place the $2k$ ordered parts of $\pi^\pm(S)$ on the circle such that

- each ordered part should be placed consecutively and clockwise on the circle, and except for in the zero class, there should be no overlapping among them;
- each ordered part should be placed entirely on the extended right half, or entirely on the extended left half; if an ordered part is placed on the extended right half, then its negative should be placed on the extended left half according to (a).

Since an allowable placement of the extended right half gives an allowable placement of the extended left half, we can focus on what happens only on the extended right half. Thus, it boils down to put the k essentially different ordered parts there, with each part a choice of sign. To go further, there are two possibilities:

- (1) the zero class is empty. Then the number of allowable placements is already counted by Lemma 5.3, namely, it is $(p - 2n + 2k - 1)!! / (p - 2n - 1)!!$.
- (2) the zero class is filled. In such a case, if an ordered part with length 1 is placed in the zero class, then its negative must be placed there as well; while for an ordered part with length greater than 1, we can place itself or its negative entirely on the extended right half such that the corresponding consecutive part starts with the zero class. Thus, there are $2k - n_1(S)$ ways to use an ordered part to fill the zero class. Then it boils down to place the remaining $k - 1$ essentially different ordered parts on the right half. Since there are $(p - 1)/2 - n + 1$ empty boxes there in total, similar to the second case of Lemma 5.3, one can count that the latter number is $(p - 2n + 2k - 1)!! / (p - 2n + 1)!!$. Thus the total number is $(2k - n_1(S))(p - 2n + 2k - 1)!! / (p - 2n + 1)!!$.

Summing up the numbers in (1) and (2) gives the desired expression for $f(S)$. $\square$

Let G be any subset of Δ^+ . In accordance with (15), we denote by $\text{Stir}(G, k)$ the number of quasi-antichains of Δ^+ contained in G with length $n - k$. We also put

$$(26) \quad \text{Stir}_1(G, k) = \sum_S n_1(S),$$

where S runs over all the quasi-antichains of Δ^+ contained in G with length $n - k$.

Theorem 6.3. *Let $\mathfrak{g}$ be D_n , $n \geq 4$. Then for any subset $G \subseteq \Delta^+$, the characteristic polynomial of $\text{Shi}(G)$ is given by*

$$\chi(\text{Shi}(G), t) = \sum_{k=0}^n (-1)^k \left\{ \text{Stir}(G, n - k) \prod_{j=k}^{n-1} (t - 2j + 1) - \text{Stir}_1(G, n - k) \prod_{j=k}^{n-2} (t - 2j - 1) \right\}.$$

Proof. This follows from Lemma 3.7 and Lemma 6.2. $\square$

ACKNOWLEDGEMENTS

The research is supported by the National Science Foundation of China grant 11201261. In the process of carrying out this work, the author had more than a few conversations with Dr. S.-J. Wang about hyperplane arrangements. I would like to thank him heartily for sharing his knowledge with me.

REFERENCES

- [1] D. Armstrong and B. Rhoades, The Shi arrangement and the Ish arrangement, *Trans. Amer. Math. Soc.* 364 (2012), 1509–1528; arXiv:1009.1655v2.
- [2] C. A. Athanasiadis, Characteristic polynomials of subspace arrangements and finite fields, *Adv. Math.* 122 (1996), 193–233.
- [3] C. A. Athanasiadis, On free deformations of the braid arrangement, *European. J. Combin.* 19 (1998), 7–18.
- [4] V. Chari, R. J. Dolbin and T. Ridenour, Ideals in parabolic subalgebras of simple Lie algebras, *Symmetry in mathematics and physics*, 47–60, *Contemp. Math.* 490, Amer. Math. Soc., Providence, RI, 2009; arXiv:0809.0245v1.
- [5] P. Cellini and P. Papi, Ad-nilpotent ideals of a Borel subalgebra, *J. Algebra* 225 (2000), 130–141.
- [6] P. Cellini and P. Papi, Ad-nilpotent ideals of a Borel subalgebra II, *J. Algebra* 258 (2002), 112–121.
- [7] H. Crapo and G.-C. Rota, *On the foundations of combinatorial theory: combinatorial geometries*, preliminary edition, MIT Press, Cambridge, MA, 1970.
- [8] P. Headley, On a family of hyperplane arrangements related to affine Weyl groups, *J. Algebraic Combin.* 6 (1997), 331–338.
- [9] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [10] B. Kostant, Eigenvalues of a Laplacian and Commutative Lie subalgebras, *Topology* 3 (1965) suppl. 2, 147–159.
- [11] B. Kostant, The set of Abelian ideals of a Borel subalgebra, *Catran decompositions and Discrete Series Representations*, *Int. Math. Res. Not.* 5 (1998), 225–252.
- [12] A. Postnikov and R. Stanley, Deformations of Coxeter hyperplane arrangements, *J. Combin. Theory Ser. A*, 91 (2000), 544–597.
- [13] C. Righi, Ad-nilpotent ideals of a parabolic subalgebra, *J. Algebra* 319 (2008), 1555–1584.
- [14] C. Righi, Enumeration of ad-nilpotent ideals of parabolic subalgebras for exceptional types, arXiv:0804.2404v1.
- [15] J.-Y. Shi, The Kazhdan-Lusztig cells in certain affine Weyl groups, *Lecture Notes in Mathematics* no. 1179, Springer-Verlag, Berlin/Heidelberg/Newyork (1986).
- [16] J.-Y. Shi, Sign types corresponding to affine Weyl groups, *J. London Math. Soc.* 35 (1987), 56–74.
- [17] J.-Y. Shi, The number of $\oplus$ -sign types, *Quart. J. Math. Oxford* 48 (1997), 93–105.
- [18] R. Stanley, *Enumerative Combinatorics*, vol. 1, Second Edition, Cambridge University Press (2012).
- [19] H. Terao, Generalized exponents of a free arrangement of hyperplanes and the Shepherd-Todd-Beriskorn formula, *Invent. Math.* 63 (1981), 159–179.
- [20] T. Zaslavsky, Facing up to arrangements: face-count formulas for partitions of space by hyperplanes, *Mem. Amer. Math. Soc.* 1 (154) (1975).

(Dong) SCHOOL OF MATHEMATICS, SHANDONG UNIVERSITY, JINAN 250100, CHINA

E-mail address: chaoping@sdu.edu.cn